



Avipox[®]

Lyophisate live attenuated Fowlpox Virus strain HP-2 Vaccine

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS:

Each dose contains:

- Active substance:
A minimum of 10^{2.7} TCID₅₀ of freeze-dried live attenuated Fowlpox Virus strain HP-2.
- Other ingredients:
Lactose and nonfat dried milk powder.

INDICATIONS:

Avipox is indicated for vaccination of tukeys, layers and breeder chicken against fowlpox disease.

CONTRAINDICATIONS:

Do not vaccinate sick or stressed birds.

ADVERSE REACTIONS:

None.

If you notice any side effects inform your veterinarian.

TARGET SPECIES:

Tukeys and chickens.

DOSAGE:

Routine practice: Vaccine administration by wing-web method between the 7th and 14th week of age.

VACCINATION SCHEDULE:

Via wing-web method:

- Practice in endemic areas: Vaccine administration from the 4th week of age, with booster vaccination 3 months later.
- Dissolve the freeze dried pellet vaccine with the cold and sterile diluent provided.
- Chicken vaccination: Wing-web method: after dipping the vaccination needle into the vial containing the diluted vaccine, puncture the unfledged skin of the wing.
- Turkey vaccination: after dipping the vaccination needle into the vial containing the diluted vaccine, puncture the skin of the middle of the thigh.

ADVICE ON CORRECT ADMINISTRATION:

- Apply usual aseptic procedures.
- Use only clean, antiseptic or disinfectant-free materials for the preparation of the vaccine solution.

WITHDRAWAL PERIOD:

Zero days.

SPECIAL STORAGE PRECAUTIONS:

- Store in a refrigerator (2°C-8°C).
- Protect from direct sunlight.
- Keep out of the reach and sight of children.
- The vaccine remains stable for 2 years from production date stated on the label under the recommended storage conditions.
- Shelf life after reconstitution: Use the vaccine within 2 hours after reconstitution.

SPECIAL WARNING(S):

Special warnings for each target species:

Vaccinate healthy birds only.

Special precautions for use in animals:

Not applicable.

Special precautions to be taken by the person administering the medicinal product to animals:

Care should be taken to avoid accidental injection.

Use during lay, fertility:

Can be administered during the laying period.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before and after any other veterinary medicinal product therefore needs to be made on a case-by-case basis.

Overdose:

10X of the dose of a vaccine: no reaction has been reported.

If you notice any side effects inform your veterinarian.

SPECIAL PRECAUTIONS FOR DISPOSAL:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

OTHER INFORMATION:

Fowlpox is a common disease of chickens, turkeys, pheasants, as well as pets and wild birds. It is caused by numerous species of the genus Avipox virus, family poxviridae. Fowlpox may occur in two forms, cutaneous or diphtheric or both. Control of fowlpox is based on immunization of birds with attenuated vaccines.

Package sizes:

Boxes of different presentations:

- Vials of: 50 doses, 100 doses, 200 doses, 500 doses, 1000 doses.
- Diluent vial.
- Connector and applicator.

FOR VETERINARY USE ONLY.



Jovac

MANUFACTURING AUTHORIZATION HOLDER AND MANUFACTURED BY:

Tel. +962 6 5232162, Fax: +962 6 5232210

P.O. Box 43, Amman 11941- Jordan

E-mail: info@jovaccenter.com, www.jovaccenter.com

06INE023.07
03/04/2023