



Jova Zeit 1,2,4[®]

Inactivated oil emulsion for injection Newcastle Disease Virus, Infectious Bronchitis Disease Virus and EDS'76 vaccine

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS:

Each dose contains:

- Active substance:
 - A minimum of 10^{8.6} EID₅₀ of inactivated Newcastle Disease Virus strain La Sota.
 - A minimum of 10^{6.8} EID₅₀ of inactivated Infectious Bronchitis Disease Virus strain M41.
 - A minimum of 1000 HA units of inactivated Adenovirus EDS'76.
- Other ingredients:
Span 80, Tween 80, Thiomersal and oil adjuvant.

INDICATIONS:

To induce active immunity in chickens against Newcastle Disease Virus, Infectious Bronchitis Disease Virus and Egg drop syndrome '76.

CONTRAINDICATIONS:

Do not vaccinate sick or stressed birds.

ADVERSE REACTIONS:

Very rare: minor reaction may occur. If you notice any side effects, please inform your veterinarian.

TARGET SPECIES:

Chickens.

DOSAGE:

- Layers:
Vaccinate 0.3 ml subcutaneously in the lower part in the back of the neck or intramuscularly in the breast or thigh muscle at 15-16 weeks of age.
- For optimal titer results, birds should be primed with live vaccine against Newcastle Disease Virus and Infectious Bronchitis Disease Virus.

ADVICE ON CORRECT ADMINISTRATION:

- Apply usual aseptic procedures.
- Bring the vaccine to room temperature.
- Shake the container well before use.

WITHDRAWAL PERIOD:

Zero days.

SPECIAL STORAGE PRECAUTIONS:

- Store in a refrigerator (2°C-8°C).
- Protect from direct sunlight.
- Do not freeze.
- Keep out of the reach and sight of children.
- The vaccine remains stable for 2 years from production date stated on the label under the recommended storage conditions.
- Shelf life after first opening: Use the vaccine within 8 hours after first opening.

SPECIAL WARNING(S):

special warnings for each target species:

Vaccinate healthy birds only.

Special precautions for use in animals:
Not applicable.

Special precautions to be taken by the person administering the medicinal product to animals:

If the vaccine is accidentally injected to the operator, urgent medical attention is required.

Use during lay, fertility:

Do not use during the laying period.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before and after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

Rare: minor reaction may occur when administering 2 X of a vaccine dose. If you notice any side effects please inform your veterinarian.

SPECIAL PRECAUTIONS FOR DISPOSAL:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

OTHER INFORMATION:

Polyvalent Inactivated oil emulsion vaccine. The viruses are grown in (SPF) chicken embryos and (SPF) duck embryos, and then inactivated and suspended in aqueous phase mixed in oil adjuvant to produce a water-in-oil emulsion vaccine.

Package sizes:

Bottles of: 300 ml (1000 doses).

FOR VETERINARY USE ONLY.



Jovac

MANUFACTURING AUTHORIZATION HOLDER AND MANUFACTURED BY:

Tel. +962 6 5232162, Fax: +962 6 5232210

P.O. Box 43, Amman 11941- Jordan

E-mail: info@jovaccen.com, www.jovaccen.com

EEINE021.07
12/09/2021